

What preventive care services are covered?

In order to prevent illness and disease, as well as minimizing health care costs, all Elevanta Health plans, including the Minimum Essential Coverage (MEC) plans, cover a list of preventative services at no out-of-pocket cost to members. That means no copayments, coinsurance or deductibles. This applies only when these services are delivered by an in-network provider.

Preventative Care Services covered 100% include:

Adults

- 1. Abdominal Aortic Aneurysm one-time screening for men of specified ages who have ever smoked
- 2. Alcohol Misuse screening and counseling
- 3. Aspirin use to prevent cardiovascular disease for men and women of certain ages
- 4. Blood Pressure Screening for all adults
- 5. Cholesterol Screening for adults of certain ages or at higher risk
- 6. Colorectal Cancer Screening for adults over 50
- 7. Depression Screening for adults
- 8. Diabetes (Type 2) Screening for adults with high blood pressure
- 9. Diet Counseling for adults at higher risk for chronic disease
- 10. **Falls Prevention** (with exercise or physical therapy and vitamin D use) for adults 65 years and over, living in a community setting
- 11. **Hepatitis B Screening** for people at high risk, including people from countries with 2% or more Hepatitis B prevalence, and U.S. born people not vaccinated as infants and with at least one parent born in a region with 8% or more Hepatitis B prevalence.
- 12. Hepatitis C Screening for adults at increased risk, and one time for everyone born 1945 1965
- 13. HIV screening for everyone ages 15 to 65, and other ages at increasedrisk
- 14. Immunization vaccines for adults (Hepatitis A & B, Herpes Zoster, Human Papillomavirus, Influenza (Flu Shot), Measles, Mumps, Rubella, Meningococcal, Pneumococcal, Tetanus, Diphtheria, Pertussis, Varicella)
- 15. Lung Cancer Screening for adults 55 80 at high risk for lung cancer because they're heavy smokers or have quit in the past 15 years
- 16. **Obesity** screening and counseling for all adults
- 17. Sexually Transmitted Infection (STI) prevention counseling for adults at higherrisk
- 18. Statin Preventative Medication for adults 40 to 75 at high risk
- 19. Syphilis screening for all adults at higher risk
- 20. Tobacco Use Screening for all adults and cessation interventions for tobaccousers
- 21. Tuberculosis Screening for certain adults without symptoms at high risk



Women, including pregnant women

- 1. Anemia Screening on a routine basis for pregnant women
- 2. **Breastfeeding** comprehensive support and counseling from trained providers, and access to breastfeeding supplies, for pregnant and nursing women
- 3. **Contraception:** Food and Drug Administration-approved contraceptive methods, sterilization procedures, and patient education and counseling, as prescribed by a health care provider for women with reproductive capacity (not including abortifacient drugs). This does not apply to health plans sponsored by certain exempt "religious employers".
 - Barrier methods, like diaphrams and sponges
 - Hormonal methods, like birth control pills and vaginal rings
 - Implanted devices, like intrauterine devides (IUDs)
 - Emergency contraception, like Plan B [®] and ella [®]
 - Sterilization procedures
 - Patient education and counseling

* Plans **are not** required to cover drugs to induce abortions and services for male reproductive capacity like, like vasectomies.

- 4. Folic Acid Supplements for women who may become pregnant
- 5. **Gestational Diabetes Screening** for women 24 to 28 weeks pregnant and those at high risk of developing gestational diabetes
- 6. **Diabetes Screening** for women with a history of gestational diabetes who aren't currently pregnant and who haven't been diagnosed with type 2 diabetes before
- 7. Gonorrhea screening for all women at higher risk
- 8. Hepatitis B screening for pregnant women at their first prenatal visit
- 9. Preeclampsia Prevention and Screening for pregnant women with high blood pressure
- 10. Rh Incompatibility screening for all pregnant women and follow-up testing for women at higher risk
- 11. Syphilis screening for all pregnant women or other women at increased risk
- 12. Expanded Tobacco Intervention and Counseling for pregnant tobacco users
- 13. Urinary tract or other infection screening for pregnant women
- 14. **Breast Cancer Genetic Test Counseling (BRCA)** risk assessment and genetic testing for women who have family members with breast, ovarian, tubal, or peritoneal cancer
- 15. Breast Cancer Mammography screenings every 1 to 2 years for women over 40
- 16. Breast Cancer Chemoprevention counseling for women at higher risk
- 17. Breast cancer risk-reducing medications, such as tamoxifen or ralozifene, for women who are at increased risk for breast cancer
- 18. Cervical Cancer screening for sexually active women. Pap test (also called a Pap smear) every 3 years for women 21-65; Human Papilomavirus (HPV) DNA test with the combination of a Pap smear every 5 years for women 30-65 who don't want a Pap smear every 3 years
- 19. Chlamydia Infection screening for younger women and other women at higher risk
- 20. Domestic and interpersonal violence screening and counseling for all women



- 21. Osteoporosis screening for women over age 60 depending on risk factors
- 22. Urinary incontinence screening for women yearly
- 23. Well-woman visits to get recommended services for women under 65

Children

- 1. Alcohol and Drug Use assessments for adolescents
- 2. Autism screening for children at 18 and 24 months
- 3. **Behavioral assessments** for children at the following ages: 0–11 months, 1–4 years, 5–10 years, 11–14 years, 15–17 years.
- 4. Bilirubin Concentration Screening for newborns
- 5. **Blood Pressure** screening for children at the following ages: es: 0–11 months, 1–4 years, 5–10 years, 11–14 years, 15–17 years.
- 6. Blood Screening for newborns
- 7. Cervical Dysplasia screening for sexually active females
- 8. **Depression** screening for adolescents
- 9. Developmental screening for children under age 3
- 10. **Dyslipidemia** screening for children at higher risk of lipid disorders at the following ages: 1–4 years, 5–10 years, 11–14 years, 15–17 years.
- 11. Fluoride Chemoprevention supplements for children without fluoride in their water source
- 12. Fluoride Varnish for all infants and children as soon as teeth are present
- 13. Gonorrhea preventive medication for the eyes of all newborns
- 14. **Hearing** screening for all newborns; and for children once between 11 and 14 years, once between 15 and 17 years, and once between 18 and 21 years
- 15. **Height, Weight and Body Mass Index** measurements for children at the following ages: es: 0– 11 months, 1–4 years, 5–10 years, 11–14 years, 15–17 years.
- 16. Hematocrit or Hemoglobin screening for children
- 17. Hemoglobinopathies or sickle cell screening for newborns
- Hepatitis B Screening for adolescents at high risk, including adolescents from countries with 2% or more Hepatitis B prevalence, and U.S.–born adolescents not vaccinated as infants and with at least one parent born in a region with 8% or more Hepatitis B prevalence: 11–17 years
- 19. HIV screening for adolescents at higher risk
- 20. Hypothyroidism screening for newborns
- 21. Immunization Vaccines for children from birth to age 18 doses, recommended ages, and recommended populations vary: <u>Diphtheria</u>, <u>Tetanus</u>, <u>Pertussis (Whooping Cough)</u>; <u>Haemophilus</u> influenza type b; <u>Hepatitis A</u>; <u>Hepatitis B</u>; <u>Human Papillomavirus (HPV)</u>; <u>Inactivated Poliovirus</u>; <u>Influenza (flu shot)</u>; <u>Measles</u>; <u>Meningococcal</u>; <u>Pneumococcal</u>; <u>Rotavirus</u>; <u>Varicella (Chickenpox)</u>



- 22. Iron supplements for children ages 6 to 12 months at risk for anemia
- 23. Lead screening for children at risk of exposure
- 24. Maternal Depression Screening for mothers of infants at 1, 2, 4, and 6-month visits
- 25. **Medical History** for all children throughout development at the following ages: es: 0–11 months, 1–4 years, 5–10 years, 11–14 years, 15–17 years.
- 26. **Obesity** screening and counseling
- 27. **Oral Health risk** assessment for young children Ages: es: 0–11 months, 1–4 years, 5–10 years.
- 28. Phenylketonuria (PKU) screening for this genetic disorder in newborns
- 29. Sexually Transmitted Infection (STI) prevention counseling and screening for adolescents at higher risk
- 30. **Tuberculin** testing for children at higher risk of tuberculosis at the following ages: 0 to 11 months, 1 to 4 years, 5 to 10 years, 11 to 14 years, 15 to 17 years.
- 31. Vision screening for all children.